



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**22.09.2004 Bulletin 2004/39**

(51) Int Cl.7: **A01N 25/18, A01M 1/20**

(21) Application number: **04101052.1**

(22) Date of filing: **15.03.2004**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IT LI LU MC NL PL PT RO SE SI SK TR**  
Designated Extension States:  
**AL LT LV MK**

- **Ogawa, Kinya, c/o SHIN-ETSU CHEMICAL CO., LTD.  
Tokyo (JP)**
- **Aiba, Noboru, c/o SHIN-ETSU CHEMICAL CO., LTD.  
Naka Kubiki-gun, Niigata-ken, Niigata (JP)**
- **Fukumoto, Takehiko, SHIN-ETSU CHEMICAL CO., LTD.  
Naka Kubiki-gun, Niigata-ken, Niigata (JP)**

(30) Priority: **17.03.2003 JP 2003071168**

(71) Applicant: **Shin-Etsu Chemical Co., Ltd.  
Chiyoda-ku, Tokyo (JP)**

(74) Representative: **Jorio, Paolo, Dr. Ing. et al  
Studio Torta S.r.l.,  
Via Viotti, 9  
10121 Torino (IT)**

(72) Inventors:  
• **Hojo, Tatsuya, c/o SHIN-ETSU CHEMICAL CO., LTD.  
Naka Kubiki-gun, Niigata-ken, Niigata (JP)**

(54) **Sustained release dispenser comprising two or more sex pheromone substances and a pest control method**

(57) Provided are a sustained release dispenser which can control the release of sex pheromone simultaneously toward two or more species of insect pests and can also release the sex pheromone in response to the generation period of insect pest and the pest control method toward two or more insect pests. Provided specifically is a sex pheromone sustained release dispenser comprising two or more sex pheromone substances, each substance being aliphatic derivative having 10 to 20 carbons, and first and second chambers wherein some or all of the sex pheromone substance having a smallest carbon number excluding a carbon number in a functional group is contained in a first polymer chamber and the remainder is contained in a second polymer chamber. Provided also is a pest control method for using the sex pheromone sustained release dispenser.

**FIG.1 (a)**




FIG.1 (b)




FIG.1 (c)





FIG.1 (d)



**Description**BACKGROUND OF THE INVENTION

## 5 1. Field of the invention

**[0001]** The invention relates to a sustained release dispenser as well as a pest control method, enabling a simultaneous release of two or more sex pheromone substances.

## 10 2. Description of the related art

**[0002]** As an effective method for utilizing a sex pheromone of an insect pest, the method has progressed to a practical stage, where a certain concentration of chemically synthesized sex pheromone is kept in a field so as to disrupt the mating of the insect pest. The important issues for the pest control method are development of the sustained release formulation (or dispenser) which can discharge a synthesized sex pheromone substance of an insect pest at a certain rate or higher in a long period; and the period of the mating disruption toward a target insect pest.

**[0003]** There are many cases where it is relatively easy to control the release of one or more sex pheromone substances of a single species of insect pest at a certain rate in a certain period because said one or more sex pheromone substances are only one kind or two or more substances having similar chemical structures. However, it is difficult to enclose sex pheromone substances with very different chemical structures in one dispenser and then control the release of the substances toward an insect pest having the substances. For instance, a diamondback moth (*Plutella xylostella*), an insect harmful to vegetables, has the sex pheromone substances (Z)-11-hexadecenal and (Z)-hexadecenyl acetate; and the beet armyworm (*Spodoptera exigua* (Hubner)(*Lepidoptera* : *Noctuidae*)), an insect harmful to vegetables, has the sex pheromone substances (Z, E)-9, 12-tetradecenyl acetate and (Z)-9-tetradecenol. Thus, each harmful insect has sex pheromone substances with different functional groups. Further, a principal component in the sex pheromone substance of leaf rollers, insects harmful to fruit trees and tea, is (Z)-11-tetradecenyl acetate and some of minor components therein are (Z)-9-dodecenyl acetate and 11-dodecenyl acetate. Thus, although the components have same functional groups, they have different carbon numbers. Hence, when the components are enclosed in a same dispenser, it is not possible to control the release of each component because of each component having different vapor pressure. It should be noted that there is rarely a single kind of insect pest for a certain crop. Thus, it is usually necessary to control two or more insect pests simultaneously in an area having various insect pests such as Japan.

**[0004]** The difficulties encountered when designing a sustained release dispenser which releases compounds having different chemical structures simultaneously are described below:

- 35 (1) The release of compounds cannot be controlled because of each compound having a different boiling point (vapor pressure) due to a different functional group thereof;
- (2) The release of compounds cannot be controlled because of each compound having a different boiling point (vapor pressure) due to a different carbon number thereof, even when the functional group of each compound is identical;
- 40 (3) A dispenser comprising a porous support without a vapor barrier will release a compound having a higher vapor pressure faster even if an evaporation area is identical.

**[0005]** It is also difficult to design a sustained release dispenser toward two or more insects at the same because the generation period of each insect pest differs.

45

SUMMARY OF THE INVENTION

**[0006]** Thus, there are cases where sex pheromone substances having different chemical structures are mixed toward a single kind of insect pest or sex pheromone substances are mixed toward two or more insect pests. However, it is difficult to release the compounds in the same ratio as the composition ratio in the dispenser. It is also difficult to control the release of the compounds in response to the generation of each kind of insect pests.

**[0007]** For instance, it is generally believed to be necessary to control three kinds of insect pests, leaf rollers, *Grapholita molesta* and *Carposina niponensis* simultaneously for the pear in Japan. The principal component of sex pheromone substances for the leaf rollers is Z-11-tetradecenyl acetate having 14 carbons excluding the carbons of the ester group. The principal component of the sex pheromone substances of the oriental fruit moth (*Grapholita molesta*) is Z-8-dodecenyl acetate having 12 carbons excluding the carbons of the ester group. The principal component of the sex pheromone substances of the peach fruit moth (*Carposina niponensis*) is Z-13-icosen-10-one having 20 carbons excluding the carbon of the carbonyl group. Although the vapor pressure of Z-8-dodecenyl acetate of the *Grapholita*

*molesta* sex pheromone substance is higher than the vapor pressures of sex pheromone substances of the other two insect pests, a long-term release of Z-8-dodecenyl acetate is desirable. It is because it is important to protect the late-maturing cultivars of autumn from the insect pests. A sustained release dispenser which can satisfy the above requirement has not yet been developed.

5 [0008] On the other hand, in order to disrupt the mating of two or more species of insect pests, it is considered to design a dispenser for each type of insect pest sex pheromone substance and then place each individual dispenser to release the sex pheromone substance at different period in response to the generation of the insect. However, it requires high cost for the preparation of dispensers and large labor for the placements of dispensers. Consequently, it opposes to the present trend of laborsaving agriculture.

10 [0009] Thus, the development of a laborsaving sustained release dispenser which can release two or more sex pheromone substances in a desired manner has been anticipated.

[0010] The purpose of this invention is to provide a sustained release dispenser which can control the simultaneous release sex pheromone substances in case where the sex pheromone substances have different chemical structures of a single species of insect pest are mixed, and/or where the sex pheromone substances of two or more species of insect pests are mixed. In the latter case, the sustained release dispenser which can release the sex pheromone substances in response to the generation period of the insect pests are provided together with a pest control method.

15 [0011] The inventors studied for the sex pheromone dispensers which can solve the above problems, can control the simultaneous release of sex pheromone substances, and can release the sex pheromone substances in response to the generation period of the insect pests. Consequently, the inventors have found that a sustained release dispenser comprising some or all of the sex pheromone substance having the lowest carbon number excluding the carbon number of the functional group in one tube of polymer material, and the remainder of the sex pheromone substances in the other tube of polymer material is suitable for said purpose. It is preferable to bond these two tubes together in a parallel way and have them function as one unit. Then the invention is completed.

20 [0012] According to the present invention, a sex pheromone substance sustained release dispenser comprising at least two sex pheromone substances, each being aliphatic derivative having 10 to 20 carbons, and comprising first and second chambers of polymer materials wherein some or all of the sex pheromone substance having a lowest carbon number is in the first chamber and the remainder of the sex pheromone substances is in the second chamber. Also provided is a pest control method of using the sex pheromone substance sustained release dispenser. In addition, provided is a container for a sex pheromone substance sustained release dispenser comprising at least two sex pheromone substances of aliphatic derivatives each having 10 to 20 carbons and comprising first and second chambers of polymer materials, wherein some or all of the sex pheromone substance having a lowest carbon number is placed in the first chamber and the remainder of the sex pheromone substances is in the second chamber.

25 [0013] When the sustained release dispenser and the pest control method of this invention are used, it is possible to release sex pheromone substances simultaneously to control two or more insect pests, and to release sex pheromone substances in response to the generation periods of the insect pests.

#### BRIEF DESCRIPTION OF THE DRAWING

40 [0014]

Figure 1 illustrates the process of the sustained release dispenser of this invention, showing two tubes made of polymer material in Figure 1(a), a joint in Figure 1(b), the sealing process in which a heat sealing device is used in Figure 1(c) and the cross section of the sustained release dispenser in Figure (d).

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

45 [0015] The invention will be explained in more detail below.

[0016] An example of sex pheromones toward insect pests is shown in Table 1. Chemical structure, relation with a target insect and carbon number of each component are listed in the pest control of the pear and peach in Japan. The vapor pressure was measured by the gas flow method at 25°C according to the "Addendum to the OECD chemical test guideline, issues 7 and 8" (Adopted on July 27, 1995)..

Table 1

Type of compound	Name of compound	Carbon number of Skelton	Biological activity	Vapor pressure (20°C) (Pa)
Alcohol	Z-8-dodecene-1-ol	12	Minor component Of Grapholita molesta pheromone	0.980
	Z-11-tetradecene-1-ol	14	Minor component Of leaf roller Pherone	0.070
Ester	Z-8-dodecenyl acetate	12	Main component of Grapholita molesta	0.267
	Z-9-dodecenyl acetate	12	Minor component of leaf roller pheromone	0.267
	11-dodeceny acetate	12	Minor component of leaf roller pheromone	0.246
	10-methyldodecyl acetate	13	Minor component of leaf roller pheromone	0.186
	Z-11-tetradecenyl acetate	14	Main component of leaf roller pheromone	0.100
	Z-9-tetradecenyl acetate	14	Minor component of leaf roller pheromone	0.100
ketone	Z-13-icosen-10-one	20	Component of Carposina niponensis pheromone	0.005

**[0017]** The above compounds are contained in a narrow tube, a capsule or a laminate of polymer material membrane and the release of the compounds are controlled by a barrier of the membrane. In this case, (1) vapor pressure of the enclosed sex pheromone substance and (2) affinity (solubility parameter) between the functional group of sex pheromone substance and the polymer material become important factors in the release control.

**[0018]** Because minor components in Table 1 are of small content, the main components are focused. It is considered that the difference in the affinity between each ester and the polymer material is extremely small and the difference between each ester and ketone is also relatively small. Thus, when a uniform solution containing all compounds in Table 1 is enclosed in a certain polymer material, the release rate of each component greatly depends on its vapor pressure. That is, esters having a higher vapor pressure such as 11-dodecenyl acetate, Z-8-dodecenyl acetate and Z-9-dodecenyl acetate will be released faster. Each of these compounds has 12 carbons excluding the carbon number of the functional group. On the other hand, compounds having a lower vapor pressure such as Z-11-tetradecenyl acetate and Z-13-icosen-10-one will be released slowly. Compounds which will be released faster cannot be released uniformly so that they result in shorter life time (effective release period) of a sustained release dispenser.

**[0019]** Hence, the inventors conducted various examinations and created a sustained release dispenser effective in cases when sex pheromone substances of a single species of insect pest having different chemical structures are mixed and/or when sex pheromone substances of two or more species of insect pests are mixed. The sustained release dispenser comprises first and second chambers of polymer materials, wherein some or all of mixture of the sex pheromone substances having a lowest carbon number (excluding the carbon number of the functional group) and having a high vapor pressure such as Z-8-dodecenyl acetate and Z-8-dodecene-1-ol in Table 1 is enclosed in the first chamber and the remainder of the sex pheromone substances is enclosed in the second chamber. It is preferable to bond the first and second chambers together in a parallel way so that they form one unit. Consequently, the pheromones do not

suppress the release of the other pheromones. Therefore, the sex pheromone component of *Grapholita molesta* having the highest vapor pressure and the sex pheromone components of leaf rollers and *Carposina niponensis* can be released simultaneously in a long period. The invention is completed by this finding.

[0020] The sex pheromone substance used in the invention is an aliphatic derivative having 10 to 20 carbons. It preferably includes aliphatic linear alcohol of lepidopterans, and ester and ketone derived from the alcohol. Two or more of the sex pheromone substances are released simultaneously. The aliphatic derivative is preferably selected from the group consisting of an aliphatic linear alcohol having 10 to 18 carbons, an aliphatic linear acetate having 12 to 20 carbons and an aliphatic linear ketone having 10 to 20 carbons. It is noted that the carbon number of the functional group of the aliphatic linear acetate or aliphatic linear ketone is 1 or 2, respectively. Hence, when the carbon number of the functional group is excluded, the aliphatic linear acetate has 10 to 18 carbons and the aliphatic linear ketone has 9 to 19 carbons.

[0021] Two or more types of sex pheromone substances of the aliphatic derivative may be two or more sex pheromone substances in the category or two or more sex pheromone substances in the different categories. A stabilizer such as an antioxidant or UV absorber, or a colorant can be contained by 10% by weight or less in the sex pheromone substances.

[0022] As shown in Table 2, three species of insect pests are targeted in the case of the mating disruption agent toward insect pests of the pear and peach in Japan.

[0023] The *Grapholita molesta* have sex pheromone containing a main component of Z-8-dodecenyl acetate and a minor component of Z-8-dodecen-1-ol, and grows from the end of April to the end of September. The leaf rollers have sex pheromone containing a main component of Z-11-tetradecenyl acetate and minor components of Z-9-tetradecenyl acetate, Z-9-dodecenyl acetate, 11-dodecenyl acetate, 10-methyl dodecyl acetate and Z-11-tetradecen-1-ol, and grow from the middle of May to the beginning of September. The *Carposina niponensis* has sex pheromone containing a component of Z-13-icosen-10-one and grows from late May to the middle of September.

[0024] It should be noted that the main component refers to the highest content of sex pheromone component. The minor component refers to sex pheromone component or components other than the main component.

Table 2

Name of insect pest	Growing period	Components of sex pheromone	Carbon carbon number of skeleton	Vapor pressure (20°C) (Pa)
Grapholita Molesta	End of April to end of September	Z-8-dodecene-1-ol	12	0.980
		Z-8-dodecenyl acetate*	12	0.267
Leaf roller	Middle of May to beginning of September	Z-9-dodecenyl acetate	12	0.267
		11-dodecenyl acetate	12	0.246
		10-methyl dodecyl acetate	13	0.186
		Z-11-tetradecenyl acetate*	14	0.100
		Z-9-tetradecenyl acetate	14	0.100
		Z-11-tetradecene-1-ol	14	0.070
Carposina niponensis	End of May to middle of September	Z-13-icosen-10-on*	20	0.005

\*The main component of natural sex pheromone for each insect pest is shown.

[0025] Some or all of the main component of the sex pheromone of *Grapholita molesta* which has the lowest carbon number is enclosed in a first independent polymer chamber, and the sex pheromones of *Carposina niponensis* and a leaf roller are enclosed in a second polymer chamber. A minor component of the sex pheromone of *Grapholita molesta* which has the same carbon number as the main component thereof may be enclosed in the second chamber. However,

the minor component of the sex pheromone of *Grapholita molesta* is preferably enclosed in the second chamber together with the main component thereof because biological functions of minor components are often unknown.

[0026] The amount of sex pheromone substance or substances enclosed in each chamber is variable depending on the release period of the sustained release dispenser, the volatility of the sex pheromone substances, and the affinity or compatibility between the substances and polymer material for the container. The amount in the first chamber may be preferably 50 to 150 mg, more preferably 100 mg, and that in the second chamber may be preferably 200 to 300 mg, more preferably 230 mg.

[0027] The polymer material used in the invention may include polyolefin, acrylic, polyester, polyamide, methacrylic and copolymer of olefin and vinyl alcohol ester. When the membrane of the polymer material is used as a barrier, tube extruded in a general method, capsule, bag, and laminate may be included. Plasticizer, lubricant, stabilizer or colorant may be added in the processing stage. According to the invention, the polymer materials of the chambers or containers may be identical or different as long as they can be joined. The polymer materials of the chambers or containers may be preferably integrated to be one.

[0028] The shape of chamber or container made of polymer material is not limited as long as it can contain sex pheromone substance. For example, the sex pheromone substance may be contained in a narrow tube, capsule or laminate container, each comprising polymer membrane. The membrane can function as a barrier so that the release of the substance can be controlled.

[0029] The size of the polymer membrane can be selected in relation to the property of the contained sex pheromone substance. Thickness and the inside and outside diameters of the first and second chambers may be same or different. The thickness, in particular, is greatly influenced by the evaporation rate of the sex pheromone component so that the appropriate thickness can be selected in view of a necessary release period, a vapor pressure and solubility parameter of the sex pheromone component. It should be noted it is difficult to seal or joint the chambers containing the pheromone substance when the thickness of the first polymer chamber is extremely different from that of the second polymer chamber. Thus, the thickness has to be selected in practical range.

[0030] According to the invention, the construction of the first or second polymer chamber is not particularly limited. However, the construction which can be chemically connected by heat seal or adhesive may be preferable. Physical connection with tape or wire may be possible. Tube 1 of a first polymer chamber and tube 2 of a second polymer chamber may be separate as shown in Figure 1(a), or joined at joint 3 in a parallel manner as shown in Figure 1(b). It may be possible to connect the ends of two tubes together chemically or physically by using heat seal device 4 as shown in Figure 1(c) so as to produce the dispenser comprising sex pheromones A1 and A2 as shown in the cross section of Figure 1(d). It should be noted that the invention is not limited to the chamber constructions shown in figures.

[0031] This invention will be explained in detail in the following Examples and Comparative Examples, the use of which, however, is not limited to the following examples.

#### Example 1

[0032] Sustained release dispenser for controlling three insect species: *Grapholita molesta*, *Carposina niponensis* and leaf roller.

[0033] Dispensers in the form of 20-cm-long tubes (A) and (B) joined in a parallel manner as shown in Figure 1 were prepared. The contents of tubes (A) and (B) are shown below. The dispensers were placed in 2 ha of a pear orchard in Chiba Prefecture, Japan, on May 9, as shown in Table 3. The dispensers were placed in ratio of 180 tubes per 10 a. As a result, damage caused by leaf roller, *Carposina niponensis* and *Grapholita molesta* was not observed in fruits up to harvest time of mid October. Residual percentages of each pheromone component of each species of insects for the tubes placed in the orchard are shown in Figure 3.

(A) The following components were contained in a high-density polyethylene tube with inside diameter of 1.4 mm and outside diameter of 2.5 mm (tube thickness of 0.55 mm):

#### Leaf roller components

Z-11-tetradecenyl acetate: 88 mg  
 Z-9-tetradecenyl acetate: 17 mg  
 10-methyldodecyl acetate: 2 mg  
 Z-9-dodecenyl acetate: 5 mg  
 11-dodecenyl acetate: 2 mg  
 Z-11-tetradecene-1-ol: 1 mg

#### *Carposina niponensis* component

EP 1 459 626 A1

Z-13-icosane-10-one : 80 mg

*Grapholita molesta* components

Z-8-dodecenyl acetate: 34 mg  
Z-8-dodecene-1-ol: 0.3 mg

(B) The following components were contained in a high-density polyethylene tube with inside diameter of 0.90 mm and outside diameter of 2.30 mm (tube thickness of 0.70 mm)

*Grapholita molesta* components

Z-8-dodecenyl acetate: 95 mg  
Z-8-dodecene-1-ol: 1 mg

Table 3

Date	Elapsed days (day)	Residual percentage (%)		
		<i>Grapholita molesta</i>	Leaf roller	<i>Carposina niponensis</i>
May 9	0	100	100	100
Jun.8	30	68.9	85.7	94.7
Jul.6	58	45.9	71.2	88.3
Aug.8	91	16.1	45.2	73.9
Sep.6	120	8.2	26.3	53.4
Oct.6	150	4.8	17.2	44.1

Comparative Example 1

**[0034]** A 20-cm-long high-density polyethylene tube with inside diameter of 1.5 mm and outside diameter of 2.6 mm (thickness of 0.55 mm) was bonded with aluminum wire to afford it formability. The solution wherein all of the sex pheromone components used in Example 1 had been mixed uniformly was contained in the tube so as to produce the dispenser. Some of the dispensers were placed in said pear orchard. Changes of residual percentage of each pheromone component over elapsed time were measured.

**[0035]** The results are shown in Table 4 below.

Table 4

Date	Elapsed Days (day)	Residual percentage (%)		
		<i>Grapholita molesta</i>	Leaf roller	<i>Carposina niponensis</i>
May 9	0	100	100	100
Jun.8	30	51.9	89.5	96.2
Jul.6	58	25.1	79.2	89.1
Aug.8	91	2.9	55.3	80.2
Sep.6	120	2.3	45.3	75.4
Oct.6	150	1.1	27.5	59.3

**[0036]** According to the above results, the life time of the *Grapholita molesta* is believed to be about ninety days.

Example 2

**[0037]** The sex pheromone dispenser toward Peach Twig Borer (*Anarsia lineatella*) which is an insect pest of fruit trees outside of Japan comprises 20-cm-long tubes (A) and (B), joined in parallel manner as shown in Figure 1, and the sex pheromone substance. The tubes (A) and (B) are described below. Release was confirmed at 0.3 m/s in a constant temperature tank at 25°C. Changes of the residual percentage of each sex pheromone component over elapsed days are shown in Table 5.

(A) The 140 mg of E-5-dodecenyl acetate is contained in a high-density polyethylene tube with inside diameter of 1.07 mm and outside diameter of 2.07 mm (thickness of 0.50 mm).

(B) The 60 mg of E-5-dodecenol is contained in a high-density polyethylene tube with inside diameter of 0.70 mm and outside diameter of 1.10 mm (thickness of 0.20 mm).

Table 5

Elapsed days (day)	Residual percentage (%)		A/B
	Tube A E-5-dodecenyl acetate	Tube B E-5-dodecenol	
0	100	100	1.00
22	78.4	81.4	0.96
55	45.8	48.3	0.95
85	23.9	26.0	0.92
111	9.1	10.3	0.88

Comparative example 2

**[0038]** A 20-cm-long high-density polyethylene tube with inside diameter of 1.28 mm and outside diameter of 2.48 mm (thickness of 0.60 mm) was bonded with aluminum wire to afford it formability. The solution wherein all of the sex pheromone components used in Example 2 had been mixed uniformly was contained in the tube so as to produce the dispenser. The dispensers were tested in the same manner as in Example 2. Changes of residual percentage of each pheromone component over elapsed time were measured.

Table 6

Elapsed days (day)	Residual percentage (%)		A/B
	Tube A E-5-dodecenyl acetate	Tube B E-5-dodecenol	
0	100	100	1.00
22	83.2	96.0	0.87
55	49.2	75.6	0.65
85	28.7	57.9	0.50
111	16.0	44.5	0.36

**Claims**

1. A sex pheromone sustained release dispenser comprising two or more sex pheromone substances, each substance being an aliphatic derivative having 10 to 20 carbons, and first and second chambers, wherein some or all of the sex pheromone substance having a smallest carbon number excluding a carbon number in a functional group is contained in the first polymer chamber and the remainder is contained in the second polymer chamber.
2. The sex pheromone sustained release dispenser according to Claim 1 wherein said aliphatic derivative is selected

**EP 1 459 626 A1**

from the group consisting of aliphatic linear alcohol having 10 to 18 carbons, aliphatic linear acetate having 12 to 20 carbons and aliphatic linear ketone having 10 to 20 carbons.

- 5
3. The sex pheromone sustained release dispenser according to Claim 1 or 2 wherein said sex pheromone sustained release dispenser is toward two or more species of insect pests.
4. A method of pest control using said sex pheromone sustained release dispenser according to any one of Claim 1 to 3.
- 10
5. A container for a sex pheromone sustained release dispenser comprising two or more sex pheromone substances, the container comprising a first polymer chamber for receiving some or all of the sex pheromone substance having a smallest carbon number excluding a carbon number of a functional group, and a second polymer chamber for receiving the remainder.

15

20

25

30

35

40

45

50

55

FIG.1 (a)




FIG.1 (b)




FIG.1 (c)





FIG.1 (d)





European Patent  
Office

EUROPEAN SEARCH REPORT

Application Number  
EP 04 10 1052

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	US 6 419 943 B1 (SAKURADA TOYOHISA ET AL) 16 July 2002 (2002-07-16) * column 6, line 27-49 * * column 15, line 8-53; figures 1,2 * ---	1-5	A01N25/18 A01M1/20
Y	EP 0 540 932 A (BASF AG) 12 May 1993 (1993-05-12) * page 5, line 31-34; figure 2A * ---	1-5	
Y	DE 26 41 630 A (ALBANY INT CORP) 11 August 1977 (1977-08-11) * page 28, paragraph 4 - page 31, paragraph 1 * ---	1-5	
Y	US 4 600 146 A (OHNO SHIGERU) 15 July 1986 (1986-07-15) * column 6, line 56-68; figure 5 * ---	1-5	
Y	EP 0 913 088 A (SHINETSU CHEMICAL CO) 6 May 1999 (1999-05-06) * abstract; figures * ---	1-5	
Y	EP 0 342 126 A (SHINETSU CHEMICAL CO) 15 November 1989 (1989-11-15) * page 3, line 12,13 * * page 7, line 35-38 * ---	1-5	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A01N A01M
Y	EP 0 938 842 A (SHINETSU CHEMICAL CO) 1 September 1999 (1999-09-01) * column 2, line 8 - column 7, line 12 * -----	1-5	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 9 July 2004	Examiner Schlichting, N
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03 82 (P04/C01)

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 04 10 1052

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-07-2004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6419943 B1	16-07-2002	JP 10008031 A	13-01-1998
		JP 10017407 A	20-01-1998
		JP 10017408 A	20-01-1998
		AU 723108 B2	17-08-2000
		AU 2684497 A	15-01-1998
		EP 0816430 A2	07-01-1998
		US 6159489 A	12-12-2000
		US 5993843 A	30-11-1999
EP 0540932 A	12-05-1993	DE 4136212 A1	06-05-1993
		CA 2081728 A1	03-05-1993
		EP 0540932 A1	12-05-1993
		US 5316148 A	31-05-1994
DE 2641630 A	11-08-1977	US 4017030 A	12-04-1977
		AR 220507 A1	14-11-1980
		AU 508179 B2	13-03-1980
		AU 1332076 A	27-10-1977
		BR 7606411 A	31-05-1977
		CA 1064391 A1	16-10-1979
		CH 615806 A5	29-02-1980
		DE 2641630 A1	11-08-1977
		DK 434576 A	27-03-1977
		FI 761919 A ,B,	27-03-1977
		FR 2325400 A1	22-04-1977
		GR 65991 A1	13-01-1981
		IL 49511 A	13-09-1981
		IN 143834 A1	11-02-1978
		IT 1066214 B	04-03-1985
		JP 52055969 A	07-05-1977
		NL 7610606 A	29-03-1977
NO 761418 A ,B,	29-03-1977		
NZ 180698 A	11-01-1979		
SE 424595 B	02-08-1982		
US 4600146 A	15-07-1986	JP 1350282 C	28-11-1986
		JP 57156403 A	27-09-1982
		JP 61016361 B	30-04-1986
		AU 534192 B2	12-01-1984
		AU 8071482 A	30-09-1982
		CA 1189785 A1	02-07-1985
		DE 3262590 D1	25-04-1985
		EP 0061192 A1	29-09-1982
EP 0913088 A	06-05-1999	AU 702709 B1	04-03-1999
		EP 0913088 A1	06-05-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 04 10 1052

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-07-2004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0913088	A	JP 11225646 A	24-08-1999
		US 6216960 B1	17-04-2001
-----			
EP 0342126	A	15-11-1989	JP 2049702 A
			20-02-1990
			JP 2108624 C
			21-11-1996
			JP 8018927 B
			28-02-1996
			DE 68911027 D1
			13-01-1994
			DE 68911027 T2
			26-05-1994
			EP 0342126 A2
			15-11-1989
			US 4923119 A
			08-05-1990
-----			
EP 0938842	A	01-09-1999	DE 69912012 D1
			20-11-2003
			EP 0938842 A1
			01-09-1999
			ES 2207873 T3
			01-06-2004
			JP 11279011 A
			12-10-1999
			US 2001053372 A1
			20-12-2001
-----			